What are the possible side effects?

Adverse reactions of lozenges are generally mild.

If you notice any allergic reactions like skin itchiness, face or hand swelling, breathing difficulty, etc., consult your doctor immediately.

What if I miss a dose?

Use the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose.

Do not double the dose.

How to store?

- 1. Keep out of reach of children.
- 2. Store in a cool, dry place and away from direct heat and light.


The information in this leaflet is compiled by the Hospital Authority (IHA) for general educational purpose and reference only. Whilst efforts have been made to ensure accuracy, no guarantee can be provided as to the completeness, timeliness or usefulness of the information, or that the information is the most updated. HA cannot warrant that the information provided can meet your health or medical requirements. Always seek the advice of your doctor or pharmacist for any medical condition that you may have, rather than relying only on the information provided on this leaflet

S/N: 030Ev5 (10/2024)















Lozenges generally contain analgesic, anti-inflammatory, antiseptic or soothing ingredients used for sore throat or infections.

How to use?

- 1. Rinse your mouth with water.
- 2. Suck the lozenge until it melts for a better local action on your throat.
- 3. Avoid eating, drinking or rinsing your mouth immediately after using the lozenge.

Precautions

- 1. Use the medication as directed by your doctor.
- 2. If you have history of drug allergy, consult your doctor or pharmacist before using the medication.
- 3. Discard all medications that have expired or are no longer required.
- 4. Do not share the medication with other people.
- 5. Tell your doctor and pharmacist if you are pregnant, planning to become pregnant, or breastfeeding.
- 6. Please consult your doctor if symptoms persist or worsen after using the medication.
- 7. Do not chew or swallow the lozenges to maximize the local action.
- 8. Drink adequate amount of water may also help to relieve the throat irritation.
- 9. Lozenges may contain sugar.