

Precautions

8. Family members or people who had been in close contacts with patient should also seek medical advice.
9. Do not share the medication with other people.
10. Tell your doctor and pharmacist if you are pregnant, planning to become pregnant, or breastfeeding.
11. Please consult your doctor if symptoms persist or worsen after using the medication.

What are the possible side effects?

The common side effects include skin itchiness, redness and rash of the scalp, conjunctivitis (if in contact with eyes), etc.

If you notice any severe skin irritation or allergic reactions like skin itchiness, face or hand swelling, breathing difficulty, etc., consult your doctor immediately.

How to store?

1. Keep out of reach of children.
2. Store in a cool, dry place and away from direct heat and light.



The information in this leaflet is compiled by the Hospital Authority (HA) for general educational purpose and reference only. Whilst efforts have been made to ensure accuracy, no guarantee can be provided as to the completeness, timeliness or usefulness of the information, or that the information is the most updated. HA cannot warrant that the information provided can meet your health or medical requirements. Always seek the advice of your doctor or pharmacist for any medical condition that you may have, rather than relying only on the information provided on this leaflet

S/N: 037Ev6 (10/2024)

Permethrin Liquid 1% (for Head Lice)



藥劑職系及服務統籌委員會
COC-Grade (Pharmaceutical Services)

What is Permethrin Liquid?

Permethrin liquid is a liquid preparation to be applied on scalp and hair for the eradication of head lice and their eggs.



What are head lice?

Head lice are tiny parasites that only live on the human head. They are usually found on the hair, most often around the hairline behind the ears or near the neck. Their eggs are white, yellow, brown or grey in colour, usually shiny, oval in shape, firmly attached to the hair shaft and can be found near the root of the hair under careful inspection. The most common symptom is tickling and itchy sensation on the scalp. Excessive scratching may cause bleeding or sores of the scalp, increasing the risk of skin infection.

The main route of transmission is through direct head-to-head contact with infested people or through sharing of personal items. Do not share personal items such as combs, hats, hair accessories, towels or pillows.

It is advisable to maintain good personal hygiene, wash hair regularly and thoroughly, and wash bed linen and clothing frequently.



How to use?

1. Wash your hair with a shampoo that does not contain a conditioner.
2. Rinse and towel dry, leaving hair slightly damp.
3. Shake the bottle thoroughly. Apply the liquid on the hair. Rub vigorously onto the scalp and thoroughly through the hair to ensure complete coverage, especially hair root, back of your neck, sideburns, fringe and the back of your ears.
4. Leave it on for 10 minutes. Rinse it out thoroughly with warm water and dry with a towel.
5. Comb your hair thoroughly starting from the hair root with a nit-removal comb when the hair is still wet, to remove all lice and eggs attached on the hair.
6. Within 2-3 weeks after treatment, use the nit-removal comb to check the hair for live lice every 2-3 days until sure that all lice are gone.
7. Do not re-wash the hair for 1-2 days after the lice medicine is removed.
8. If repeat application is required, use the medication as directed by your doctor or follow the instructions on the drug package.



Precautions

1. For external use only. Please seek medical advice immediately if ingested accidentally.
2. Use the medication as directed by your doctor.
3. If you have history of drug allergy, consult your doctor or pharmacist before using the medicine.
4. Discard all medications that have expired or are no longer required.
5. Wear disposable gloves when applying the liquid to prevent irritation. Wash hands thoroughly after use.
6. Avoid contact with eyes. If the liquid accidentally gets into your eyes, wash immediately with plenty of water and seek medical advice if necessary.
7. All hats, towels, pillows cases, beddings and clothing used by the patient before treatment should be washed separately from those of their family members or other old age home residents in hot water (60°C or above) for not less than 10 minutes before re-use. Combs and items that are not resistant to heat can be soaked in a mixture of bleaching agent and water (in 1:1 ratio) for 30 minutes. Items that are not washable, such as pillow and mattress, can be sealed in a plastic bag and stored for 2 weeks before re-use.

